

Molecular Defects in the Ehlers-Danlos Syndrome

SHELDON R. PINNELL, M.D.

Division of Dermatology, Department of Medicine, Duke University Medical Center, Durham, North Carolina, U.S.A

Several abnormalities in collagen biosynthesis have been described in patients with Ehlers-Danlos syndrome. Examples of collagen structural mutations as well as post-translational enzymatic defects have been detected. Patients with hydroxylysine-deficient collagen disease (Ehlers-Danlos type VI) have diminished lysyl hydroxylase activity. One mutant enzyme has been characterized which is thermally labile and had an altered affinity for ascorbate. Another mutant enzyme had a normal requirement for cofactors but activity was diminished. Type VII Ehlers-Danlos syndrome is associated with altered processing of procollagen to collagen. Most often the disorder is associated with deficient procollagen aminoprotease activity. One patient appears to represent a structural mutation of pro α_2 (I) resulting in incomplete cleavage of the amino terminal propeptide. One family with x-linked Ehlers-Danlos syndrome (type V) has been described with altered lysyl oxidase activity. Other patients with this disorder have had normal lysyl oxidase activity. The ecchymotic form of Ehlers-Danlos syndrome (type IV) has defective type III collagen synthesis. Patients have been described with absent synthesis, diminished synthesis and diminished secretion.

The Ehlers-Danlos syndrome is a striking clinical syndrome with rubber-like skin, hypermobile joints, a tendency to bruise easily, and poor wound healing [1]. In general most forms of the disorder are compatible with normal longevity; a striking exception is the ecchymotic form of Ehlers-Danlos syndrome in which premature demise occurs regularly from arterial or intestinal rupture. In 1972, the first human molecular disorder of collagen, hydroxylysine-deficient collagen disease, was described in 2 sisters with Ehlers-Danlos syndrome [2]. Since then several ultrastructural and biochemical collagen defects have been described in the Ehlers-Danlos syndrome (Table). This manuscript will focus on 4 types of Ehlers-Danlos syndrome in which biochemical collagen defects are best understood: types VI, VII, V, and IV. The range of defects is representative of the complex nature of collagen biosynthesis. These appear to include structural and regulatory collagen genomic alterations as well as intracellular and extracellular post-translation enzymatic defects.

TYPE VI EHLERS-DANLOS SYNDROME: HYDROXYLYSINE-DEFICIENT COLLAGEN DISEASE

In 1972, Pinnell et al described 2 sisters with type VI Ehlers-Danlos syndrome who had marked hyperextensibility of skin and joints, severe scoliosis, and marfanoid features [2]. Levels of hydroxylysine in skin collagen were found to be less than one per molecule of collagen and levels of hydroxylysine-derived cross-links were markedly diminished [3]. The disorder is due to deficient lysyl hydroxylase activity [4] and is inherited as an autosomal recessive. Heterozygotes have intermediate activities

of lysyl hydroxylase. Two mutant enzymes have been characterized. One has an altered affinity for ascorbate and is thermally labile [5]. This apparently represents a structural mutation of the enzyme. The other enzyme was kinetically normal but the activity was markedly diminished [6]. This may represent a structural or regulatory mutation.

The relative hydroxylation of tissue collagens has been variable in this disorder [2, 7]. Although skin was markedly deficient in hydroxylysine, bone was less deficient and cartilage was normally hydroxylated. The explanation for this variability is unknown although enzymatic polymorphism has not been excluded. Indeed evidence for isozymes has been reported [8]. When cell lysates from a lysyl hydroxylase deficient cell strain were tested against substrates for types I and IV collagen, preferential activity was demonstrated against type IV collagen. In one study an almost normal hydroxylysine content in skin collagen was described in association with lysyl hydroxylase deficiency [9]. Subsequent analysis of these skin fibroblasts however has revealed normal lysyl hydroxylase activity (Murad S, Pinnell SR: unpublished observations and Steinmann B: personal communication).

The hydroxylysine content of complement component Clq was slightly low [10] or normal [11] and functional activity was unimpaired.

Therapy with large doses of ascorbate was effective in one patient with this disorder [12]. He was able to boost his urinary excretion of hydroxylysine although hydroxyproline excretion was coordinately increased. The effect appears to be due to an overall stimulation of collagen synthesis by ascorbate [13] rather than specific stimulation of lysyl hydroxylase activity. Kinetic studies of his mutant enzyme failed to detect any alteration in affinity for ascorbate [6].

TYPE VII EHLERS-DANLOS SYNDROME: ARTHROCHALASIS MULTIPLEX CONGENITA

In 1973, Lichtenstein et al reported studies on 3 type VII Ehlers-Danlos patients with a defect in conversion of procollagen to collagen [14]. These patients had short stature, hyperextensible joints, and bilateral hip dislocation. Analysis of collagen extracted from their skin revealed elongated α_1 (I) and α_2 (I) chains resulting from inefficient conversion of type I procollagen. Their cultured skin fibroblasts had deficient procollagen aminoprotease activity. Uncleaved aminoterminal propeptides apparently interfere with fibrillogenesis and intermolecular crosslinking resulting in the fragile connective tissue.

This disorder, inherited as an autosomal recessive, is biochemically similar to dermatosparaxis found in sheep [15] and cattle [16]. The striking skin fragility characteristic of dermatosparaxis however, is absent in the Ehlers-Danlos patients. The reason for this difference is not understood.

Recently another patient with type VII Ehlers-Danlos syndrome has been described with inefficient conversion of procollagen to collagen [17]. This patient has normal levels of procollagen aminoprotease. Conversion of procollagen appears to be impeded by a structural mutation in her pro α_2 (I) near the protease cleavage site. Pro α_1 (I) is normally converted but the aminopropeptide of pro α_2 (I) remains uncleaved apparently resulting in similar difficulties in fibrillogenesis found in other patients with this form of Ehlers-Danlos syndrome. This patient has, in addition to an abnormal pro α_2 (I), an equal complement of normal pro α_2 (I). She apparently represents a new structural

This work was supported by grant 5 RO1 AM-17128 from the National Institutes of Health.

This is publication number 121 from the Dermatological Research Laboratories of Duke University Medical Center.

Reprint requests to: Sheldon R. R. Pinnell, M.D., Division of Dermatology, Department of Medicine, Duke University Medical Center, Durham, NC 27710.

Ultrastructural and biochemical collagen defects in Ehlers-Danlos syndrome

Type	Major clinical features	Ultrastructural defect	Biochemical defect
I (Gravis)	Marked joint and skin hyperextensibility. Fragile skin, poor wound healing and easy bruisability.	Variable collagen fibril diameter [29-31]	Unknown
II (Mitis)	Small joint hypermobility	Variable collagen fibril diameter [30,31]	Unknown
III (Benign hypermobile)	Large joint hypermobility	Small collagen fibril diameter [30]	Unknown
IV (Ecchymotic)	Marked bruisability, arterial and intestinal rupture	Small collagen fibril diameter [27] Dilated endoplasmic reticulum [26,27]	Diminished [25-28] or absent [23] type III collagen synthesis. Some patients may have secretion defect [26,27]
V (X-linked)	Moderate joint hypermobility. Heart valve prolapse.	Variable collagen fibril diameter [30,31]	Some patients may have lysyl oxidase deficiency [17]
VI (Hydroxyllysine-deficient)	Marked joint hypermobility. Kyphoscoliosis. Ocular fragility	None detected [2]	Lysyl hydroxylase deficiency [4-6]
VII (Arthrochalasia multiplex congenita)	Marked joint hypermobility. Dislocated hips	Not determined	Procollagen aminoprotease deficiency [14,17] One patient with structural defect in pro α 1(I) [17]
VIII (Periodontitis)	Periodontitis. Skin fragility	Not determined	Unknown
VIII (Mental retardation)	Hernias. Mental retardation	Not determined	Unknown

mutation resulting in one normal and one mutant allele for pro α ₂ (I).

TYPE V EHLERS-DANLOS SYNDROME: X-LINKED

In 1975, Di Ferrante et al reported studies on a patient with type V Ehlers-Danlos syndrome whose 2 male cousins were similarly affected [18]. The mothers were sisters, thus the disease was probably inherited as an X-linked recessive. Studies of skin fibroblasts in culture revealed increased collagen solubility as well as deficient lysyl oxidase activity. It should be noted that the lysyl oxidase activity in the control cell strain was minimal and the study failed to take into account the limited solubility of lysyl oxidase [19] and the striking effect of cellular density on lysyl oxidase activity [20]. Later Siegel, Black, and Bailey examined other patients with X-linked Ehlers-Danlos and found normal levels in skin of lysyl oxidase and intermolecular collagen crosslinks [21]. Subsequently Byers et al described a family with X-linked cutis laxa associated with lysyl oxidase deficiency apparently related to abnormal copper metabolism [22]. These patients had hyperextensible joints, a clinical feature more characteristic of Ehlers-Danlos syndrome than cutis laxa.

TYPE IV EHLERS-DANLOS SYNDROME

Clinical characteristics are different from other forms of Ehlers-Danlos syndrome in type IV Ehlers-Danlos syndrome. Bruising is prominent; skin is thin, transparent and fragile; tissues are not distensible; and healing may occur with hypertrophic scarring or keloids. Rupture of large arteries or perforation of gastrointestinal structures are common and life-threatening. Although autosomal dominant and autosomal recessive forms of this disorder occur, biochemical studies have been carried out only on the autosomal recessive form.

In 1975, Pope et al reported a patient with type IV Ehlers-

Danlos syndrome with absent type III collagen synthesis [23]. Skin, aorta, gut, bone, and tendon were obtained shortly after death. Cyanogen bromide cleavage of these tissues revealed absence of peptides characteristic of type III collagen. Skin fibroblasts in culture failed to synthesize any detectable type III procollagen. In addition, characteristic cellular staining was absent in cultured skin fibroblasts from the patient using antisera specific for type III collagen and procollagen [24]. This patient may have had a gene deletion for type III procollagen.

Subsequent studies of fibroblasts from patients with type IV Ehlers-Danlos syndrome have revealed diminished but not absent levels of type III collagen synthesis [25, 26]. They may represent structural or regulatory mutations. Electron microscopic studies have revealed small collagen fiber diameters [26, 27]. In 2 patients distended endoplasmic reticulum has been demonstrated in skin [28] and lung [26] fibroblasts. These fibroblasts demonstrated diminished total collagen synthesis and markedly deficient type III collagen synthesis [26, 28]. They may represent mutations which interfere with cellular secretion.

The Ehlers-Danlos phenotype is obviously associated with defective collagen structure. The variety of structural defects and abnormalities in post-translational modifications represented by the different forms of Ehlers-Danlos syndrome are characteristic of the complicated nature of collagen biosynthesis.

REFERENCES

- Pinnell SR: The Ehlers-Danlos syndrome in Textbook of Medicine. Edited by PB Beeson, W McDermott, JB Wyngaarden, WB Saunders Co, Philadelphia, 15th edition, 1979, pp 2054-2055
- Pinnell SR, Krane SM, Kenzora JE, Glimcher MJ: A heritable disorder of connective tissue: hydroxyllysine-deficient collagen disease. *N Engl J Med* 286:1013-1020, 1972
- Eyre DR, Glimcher MJ: Reducible cross-links in hydroxyllysine-deficient collagens of a heritable disorder of connective tissue. *Proc Natl Acad Sci USA* 69:2594-2598, 1972
- Krane SM, Pinnell SR, Erbe RW: Lysyl-procollagen hydroxylase deficiency in fibroblasts from siblings with hydroxyllysine-deficient collagen. *Proc Natl Acad Sci USA* 69:2899-2903, 1972
- Quinn RS, Krane SM: Abnormal properties of collagen lysyl hydroxylase from skin fibroblasts of siblings with hydroxyllysine-deficient collagen. *J Clin Invest* 57:83-93, 1976
- Miller RL, Elsas LJ III, Priest RE: Ascorbate action on normal and mutant human lysyl hydroxylases from cultured dermal fibroblasts. *J Invest Dermatol* 72:241-247, 1979
- Ihme A, Krieg T, Risteli L, Risteli J, Glanville R, Rautenberg J, Muller PK: Heterogeneity of ED type VI. *J Invest Dermatol* 74:455, 1980
- Risteli L, Risteli J, Ihme A, Krieg T, Muller PK: Preferential hydroxylation of type IV collagen by lysyl hydroxylase from Ehlers-Danlos syndrome type VI fibroblasts. *Biochem Biophys Res Commun* 96:1778-1784, 1980
- Steinmann B, Gitzelmann R, Vogel A, Grant ME, Harwood R, Sear CHJ: Ehlers-Danlos syndrome in two siblings with deficient lysyl hydroxylase activity in cultured skin fibroblasts but only mild hydroxyllysine deficient in skin. *Helv Paediat Acta* 30:255-274, 1975
- Pinnell SR: Abnormal collagens in connective tissue diseases. *Birth Defects* XI:23-30, 1975
- Hanauske-Abel HM, Rohm KH: The collagenous part of Cl₂ is unaffected in the hydroxyllysine-deficient collagen disease. *FEBS Letters* 110:73-76, 1980
- Elsas LJ, Miller RL, Pinnell SR: Inherited human collagen lysyl hydroxylase deficiency—ascorbic acid response. *J Pediatr* 92:378-384, 1978
- Murad S, Grove D, Lindberg KA, Reynolds G, Sivarajah A, Pinnell SR: Regulation of collagen synthesis by ascorbic acid. *Proc Natl Acad Sci USA* 78:2879-2882, 1981
- Lichtenstein JR, Martin GR, Kohn LD, Byers PH, McKusick VA: Defect in conversion of procollagen to collagen in a form of the Ehlers-Danlos syndrome. *Science* 182:298-299, 1973
- Becker U, Timpl R, Helle O, Prockop DJ: NH₂-terminal extensions on skin collagen from sheep with genetic defect in conversion of procollagen into collagen. *Biochemistry* 15:2853-2862, 1976
- Lapiere CM, Lenaers A, Kohn LD: Procollagen peptidase: an enzyme excising the coordination peptides of procollagen. *Proc Natl Acad Sci USA* 68:3054-3058, 1971
- Steinmann B, Tuderman L, Peltonen L, Martin GR, McKusick V, Prockop DJ: Evidence for a structural mutation of procollagen type I in a patient with the Ehlers-Danlos syndrome type VII. *J Biol Chem* 255:8887-8893, 1980

18. Di Ferrante N, Leachman RD, Angelini P, Donnelly PV, Francis G, Almazen A: Lysyl oxidase deficiency in Ehlers-Danlos syndrome type V. *Connect Tissue Res* 33:49-53, 1975
19. Narayanan AS, Siegel RC, Martin GR: Stability and purification of lysyl oxidase. *Arch Biochem Biophys* 162:231-237, 1974
20. Gonnerman WA, Ferrera R, Franzblau C: Measurement of medium lysyl oxidase activity in aorta smooth muscle cells. Effects of multiple medium changes and inhibition of protein synthesis. *Biochemistry* 20:3864-3867, 1981
21. Siegel RC, Black CM, Bailey AJ: Cross-linking of collagen in the X-linked Ehlers-Danlos type V. *Biochem Biophys Res Commun* 88:281-287, 1979
22. Byers PH, Siegel RC, Holbrook KA, Narayanan AS, Bornstein P, Hall JC: X-linked cutis laxa. Defective cross-link formation in collagen due to decreased lysyl oxidase activity. *N Engl J Med* 303:61-65, 1980
23. Pope FM, Martin GR, Lichtenstein JR, Penttinen R, Gerson B, Rowe DW, McKusick VA: Patients with Ehlers-Danlos syndrome type IV lack type III collagen. *Proc Natl Acad Sci USA* 72:1314-1316, 1975
24. Gay S, Martin GR, Muller PK, Timpl R, Kuhn K: Simultaneous synthesis of types I and III collagen by fibroblasts in culture. *Proc Natl Acad Sci USA* 73:4037-4040, 1976
25. Aumailley M, Krieg T, Dessau W, Muller PK, Timpl R, Bricaud H: Biochemical and immunological studies of fibroblasts derived from a patient with Ehlers-Danlos syndrome type IV. Demonstrate reduced type III collagen synthesis. *Arch Dermatol Res* 269:169-177, 1980
26. Clark JG, Kuhn C III, Uitto J: Lung collagen in type IV Ehlers-Danlos syndrome: Ultrastructural and biochemical studies. *Amer Rev Resp Dis* 122:971-978, 1980
27. Holbrook KA, Byers PH: Ultrastructural characteristics of the skin in a form of the Ehlers-Danlos syndrome type IV. *Lab Invest* 44:342-350, 1981
28. Byers PH, Holbrook KA, Smith LT, Bornstein P, McGillivray B, MacLeod PM: Altered secretion of type III collagen in a form of type IV Ehlers-Danlos syndrome (EDS). *Clin Res* 26:512A, 1978
29. Vogel A, Holbrook KA, Steinmann B, Gitzelmann R, Byers PH: Abnormal collagen fibril structure in the gravis form (type I) of Ehlers-Danlos syndrome. *Lab Invest* 40:201-206, 1979
30. Sevenich M, Schultz-Ehrenburg U, Orfanos CE: Ehlers-Danlos-Syndrome: Eine fibroblasten- und kollagenkrankheit. Typisierung und elektronenmikroskopische befunde bei funf kranken. *Arch Dermatol Res* 267:237-251, 1980
31. Black CM, Gathercole LJ, Bailey AJ, Beighton P: The Ehlers-Danlos syndrome: an analysis of the structure of the collagen fibres of the skin *Br J Dermatol* 102:85-96, 1980